Global School-based Student Health Survey

Kenya (Pastoral Region)



2003 Fact Sheet

The 2003 Kenya (Pastoral Region) GSHS measured alcohol and other drug use; sexual behaviors that contribute to HIV infection, other STI, and unintended pregnancy; unintentional injuries and violence; hygiene; dietary behaviors and overweight; physical activity; tobacco use; mental health; and protective factors.

The 2003 Kenya (Pastoral Region) GSHS was a school-based survey of students in classes 7 and 8 and forms 1 and 2. A two-stage cluster sample design was used to produce data representative of all students in classes 7 and 8 and forms 1 and 2 in Kenya (Pastoral Region). At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 100%, the student response rate was 91%, and the overall response rate was 91%. A total of 988 students participated in the Kenya (Pastoral Region) GSHS. Students self-reported their responses to each question on a computer scannable answer sheet. For comparison purposes, only students aged 13-15 years are included in the analyses for this fact sheet.

Results for students aged 13-15 years	Total	Boys	Girls			
Alcohol and Other Drug Use						
Percentage of students who had at least one drink containing alcohol on one or more days during the past 30 days	13.0 ± 6.6	15.1 ± 7.9	9.8 ± 6.3			
Percentage of students who drank so much alcohol that they were really drunk one or more times during their life	19.9 ± 6.9	23.7 ± 6.7	14.4 ± 8.2			
Percentage of students who had a hang-over, felt sick, got into trouble with their family or friends, missed school, or got into fights, as a result of drinking alcohol one or more times during their life	24.4 ± 6.9	26.8 ± 8.3	21.2 ± 6.8			
Percentage of students who used drugs one or more times during their life	12.8 ± 5.2	12.1 ± 6.2	13.3 ± 7.2			
Sexual Behaviors That Contribute to HIV Infection, Other STI, and Unintended Pregnancy						
Percentage of students who have ever had sexual intercourse	27.1 ± 7.5	35.4 ± 9.6	14.9 ± 10.3			
Unintentional Injuries and Violence						
Percentage of students who were in a physical fight one or more times during the past 12 months	41.7 ± 11.1	41.2 ± 13.7	42.0 ± 11.5			
Percentage of students who were seriously injured one or more times during the past 12 months	61.2 ± 9.8	61.0 ± 13.4	61.1 ± 10.1			
Percentage of students who were bullied on one or more days during the past 30 days	52.2 ± 12.5	50.2 ± 12.6	54.4 ± 12.8			
Hygiene						
Percentage of students who never or rarely washed their hands before eating during the past 30 days	5.8 ± 2.0	6.1 ± 1.8	5.5 ± 2.7			
Percentage of students who never or rarely washed their hands after using the toilet or latrine during the past 30 days	18.5 ± 5.1	18.6 ± 6.0	18.5 ± 7.6			

2003

Global School-based Student Health Survey

Kenya (Pastoral Region)



2003 Fact Sheet

Results for students aged 13-15 years	Total	Boys	Girls		
Dietary Behaviors and Overweight					
Percentage of students who went hungry most of the time or always during the past 30 days because there was not enough food in their home	11.5 ± 4.7	11.7 ± 5.9	11.4 ± 3.7		
Physical Activity					
Percentage of students who were physically active for a total of at least 60 minutes per day on all 7 days during the past 7 days	10.3 ± 5.6	12.1 ± 5.0	8.0 ± 7.6		
Percentage of students who spent three or more hours per day during a typical or usual day sitting and watching television, playing computer games, talking with friends, or doing other sitting activities	31.5 ± 8.0	29.3 ± 8.2	34.6 ± 10.2		
Tobacco Use					
Percentage of students who smoked cigarettes on one or more days during the past 30 days	9.9 ± 3.2	11.1 ± 6.6	7.6 ± 3.8		
Percentage of students who used any other form of tobacco on one or more days during the past 30 days	9.8 ± 6.5	9.0 ± 6.7	10.4 ± 7.2		
Percentage of students who reported people smoking in their presence on one or more days during the past 7 days	54.8 ± 5.9	55.7 ± 7.7	53.7 ± 8.6		
Mental Health					
Percentage of students who felt lonely most of the time or always during the past 12 months	13.3 ± 6.1	12.8 ± 5.1	13.5 ± 9.2		
Percentage of students who seriously considered attempting suicide during the past 12 months	15.6 ± 5.6	17.0 ± 7.4	13.6 ± 5.1		
Percentage of students who have no close friends	9.1 ± 2.6	9.1 ± 3.0	9.2 ± 3.1		

2003

Global School-based Student Health Survey

Kenya (Pastoral Region)



2003 Fact Sheet

Results for students aged 13-15 years	Total	Boys	Girls		
Protective Factors					
Percentage of students who missed classes or school without permission on one or more days during the past 30 days	38.1 ± 12.3	35.6 ± 16.1	41.0 ± 10.0		
Percentage of students who reported that most of the students in their school were never or rarely kind and helpful during the past 30 days	33.0 ± 7.8	37.0 ± 10.5	27.4 ± 10.5		
Percentage of students whose parents or guardians never or rarely really knew what they were doing with their free time during the past 30 days	39.1 ± 9.8	46.9 ± 7.0	29.2 ± 11.8		

For additional information, please contact:

Dr. Ahmed Ogwell Ministry of Health, Nairobi, Kenya, ogwell@swiftkenya.com

2003 3